DRAFT v2.0

November 3, 1999

Search for the Charged Higgs Boson in the Decays of Top Quark Pairs in the $e\tau$ and $\mu\tau$ Channels at $\sqrt{s}$=1.8 TeV

T. Affolder,[21] H. Akimoto,[42] A. Akopian,[35] M. G. Albrow,[10] P. Amaral,[7] S. R. Amendolia,[31] D. Amidei,[24] J. Antos,[1] G. Apollinari,[35] T. Arisawa,[42] T. Asakawa,[40] W. Ashmanskas,[7] M. Atac,[10] P. Azzi-Bacchetta,[29] N. Bacchetta,[29] M. W. Bailey,[26] S. Bailey,[14] P. de Barbaro,[34] A. Barbaro-Galtieri,[21] V. E. Barnes,[33] B. A. Barnett,[17] M. Barone,[12] G. Bauer,[22] F. Bedeschi,[31] S. Belforte,[39] G. Bellettini,[31] J. Bellinger,[43] D. Benjamin,[9] J. Bensinger,[4] A. Beretvas,[10] J. P. Berge,[10] J. Berryhill,[7] S. Bertolucci,[12] B. Bevensee,[30] A. Bhatti,[35] C. Bigongiari,[31] M. Binkley,[10] D. Bisello,[29] R. E. Blair,[2] C. Blocker,[4] K. Bloom,[24] B. Blumenfeld,[17] B. S. Blusk,[34] A. Bocci,[31] A. Bodek,[34] W. Bokhari,[30] G. Bolla,[33] Y. Bonushkin,[5] D. Bortoletto,[33] J. Boudreau,[32] A. Brandl,[26] S. van den Brink,[17] C. Bromberg,[25] N. Bruner,[26] E. Buckley-Geer,[10] J. Budagov,[8] H. S. Budd,[34] K. Burkett,[14] G. Busetto,[29] A. Byon-Wagner,[10] K. L. Byrum,[2] M. Campbell,[24] W. Carithers,[21] J. Carlson,[24] D. Carlsmith,[43] J. Cassada,[34] A. Castro,[29] D. Cauz,[39] A. Cerri,[31] P. S. Chang,[1] P. T. Chang,[1] J. Chapman,[24] C. Chen,[30] Y. C. Chen,[1] M. -T. Cheng,[1] M. Chertok,[37] G. Chiarelli,[31] I. Chirikov-Zorin,[8] G. Chlachidze,[8] F. Chlebana,[10] L. Christofek,[16] M. L. Chu,[1] S. Cihangir,[10] C. I. Ciobanu,[27] A. G. Clark,[13] A. Connolly,[21] J. Conway,[36] J. Cooper,[10] M. Cordelli,[12] J. Guimaraes da Costa,[24] D. Costanzo,[31] J. Cranshaw,[38] D. Cronin-Hennessy,[9] R. Cropp,[23] R. Culbertson,[7] D. Dagenhart,[41] F. DeJongh,[10] S. Dell'Agnello,[12] M. Dell'Orso,[31] R. Demina,[10] L. Demortier,[35] M. Deninno,[3] P. F. Derwent,[10] T. Devlin,[36] J. R. Dittmann,[10] S. Donati,[31] J. Done,[37] T. Dorigo,[14] N. Eddy,[16] K. Einsweiler,[21] J. E. Elias,[10] E. Engels, Jr.,[32] W. Erdmann,[10] D. Errede,[16] S. Errede,[16] Q. Fan,[34] R. G. Feild,[44] C. Ferretti,[31] I. Fiori,[3]

B. Flaugher,[10] G. W. Foster,[10] M. Franklin,[14] J. Freeman,[10] J. Friedman,[22] H. Frisch,[7] Y. Fukui,[20] S. Gadomski,[23] S. Galeotti,[31] M. Gallinaro,[35] T. Gao,[30] M. Garcia-Sciveres,[21] A. F. Garfinkel,[33] P. Gatti,[29] C. Gay,[44] S. Geer,[10] P. Giromini,[12] V. Glagolev,[8] M. Gold,[26] J. Goldstein,[10] A. Gordon,[14] A. T. Goshaw,[9] Y. Gotra,[32] K. Goulianos,[35] H. Grassmann,[39] C. Green,[33] L. Groer,[36] C. Grosso-Pilcher,[7] M. Guenther,[33] G. Guillian,[24] R. S. Guo,[1] C. Haber,[21] E. Hafen,[22] S. R. Hahn,[10] C. Hall,[14] T. Handa,[15] R. Handler,[43] W. Hao,[38] F. Happacher,[12] K. Hara,[40] A. D. Hardman,[33] R. M. Harris,[10] F. Hartmann,[18] K. Hatakeyama,[35] J. Hauser,[5] J. Heinrich,[30] A. Heiss,[18] B. Hinrichsen,[23] K. D. Hoffman,[33] C. Holck,[30] L. Holloway,[16] R. Hughes,[27] J. Huston,[25] J. Huth,[14] H. Ikeda,[40] M. Incagli,[31] J. Incandela,[10] G. Introzzi,[31] J. Iwai,[42] Y. Iwata,[15] E. James,[24] H. Jensen,[10] M. Jones,[30] U. Joshi,[10] H. Kambara,[13] T. Kamon,[37] T. Kaneko,[40] K. Karr,[41] H. Kasha,[44] Y. Kato,[28] T. A. Keaffaber,[33] K. Kelley,[22] M. Kelly,[24] R. D. Kennedy,[10] R. Kephart,[10] D. Khazins,[9] T. Kikuchi,[40] M. Kirk,[4] B. J. Kim,[19] H. S. Kim,[23] S. H. Kim,[40] Y. K. Kim,[21] L. Kirsch,[4] S. Klimenko,[11] D. Knoblauch,[18] P. Koehn,[27] A. Köngeter,[18] K. Kondo,[42] J. Konigsberg,[11] K. Kordas,[23] A. Korytov,[11] E. Kovacs,[2] J. Kroll,[30] M. Kruse,[34] S. E. Kuhlmann,[2] K. Kurino,[15] T. Kuwabara,[40] A. T. Laasanen,[33] N. Lai,[7] S. Lami,[35] S. Lammel,[10] J. I. Lamoureux,[4] M. Lancaster,[21] G. Latino,[31] T. LeCompte,[2] A. M. Lee IV,[9] S. Leone,[31] J. D. Lewis,[10] M. Lindgren,[5] T. M. Liss,[16] J. B. Liu,[34] Y. C. Liu,[1] N. Lockyer,[30] M. Loreti,[29] D. Lucchesi,[29] P. Lukens,[10] S. Lusin,[43] J. Lys,[21] R. Madrak,[14] K. Maeshima,[10] P. Maksimovic,[14] M. Mangano,[31] M. Mariotti,[29] G. Martignon,[29] A. Martin,[44] J. A. J. Matthews,[26] P. Mazzanti,[3] K. S. McFarland,[34] P. McIntyre,[37] E. McKigney,[30] M. Menguzzato,[29] A. Menzione,[31] E. Meschi,[31] C. Mesropian,[35] C. Miao,[24] T. Miao,[10] R. Miller,[25] J. S. Miller,[24] H. Minato,[40] S. Miscetti,[12] M. Mishina,[20] N. Moggi,[31] E. Moore,[26] R. Moore,[24] Y. Morita,[20] A. Mukherjee,[10] T. Muller,[18] A. Munar,[31] P. Murat,[31] S. Murgia,[25] M. Musy,[39] J. Nachtman,[5] S. Nahn,[44] H. Nakada,[40] T. Nakaya,[7] I. Nakano,[15] C. Nelson,[10] D. Neuberger,[18] C. Newman-Holmes,[10] C.-Y. P. Ngan,[22] P. Nicolaidi,[39] H. Niu,[4] L. Nodulman,[2] A. Nomerotski,[11] S. H. Oh,[9] T. Ohmoto,[15] T. Ohsugi,[15] R. Oishi,[40] T. Okusawa,[28] J. Olsen,[43] C. Pagliarone,[31] F. Palmonari,[31]

R. Paoletti,[31] V. Papadimitriou,[38] S. P. Pappas,[44] A. Parri,[12] D. Partos,[4] J. Patrick,[10] G. Pauletta,[39] M. Paulini,[21] L. Pescara,[29] T. J. Phillips,[9] G. Piacentino,[31] K. T. Pitts,[10] R. Plunkett,[10] A. Pompos,[33] L. Pondrom,[43] G. Pope,[32] F. Prokoshin,[8] J. Proudfoot,[2] F. Ptohos,[12] G. Punzi,[31] K. Ragan,[23] D. Reher,[21] A. Ribon,[29] F. Rimondi,[3] L. Ristori,[31] W. J. Robertson,[9] A. Robinson,[23] T. Rodrigo,[6] S. Rolli,[41] L. Rosenson,[22] R. Roser,[10] R. Rossin,[29] W. K. Sakumoto,[34] D. Saltzberg,[5] A. Sansoni,[12] L. Santi,[39] H. Sato,[40] P. Savard,[23] P. Schlabach,[10] E. E. Schmidt,[10] M. P. Schmidt,[44] M. Schmitt,[14] L. Scodellaro,[29] A. Scott,[5] A. Scribano,[31] S. Segler,[10] S. Seidel,[26] Y. Seiya,[40] A. Semenov,[8] F. Semeria,[3] T. Shah,[22] M. D. Shapiro,[21] P. F. Shepard,[32] T. Shibayama,[40] M. Shimojima,[40] M. Shochet,[7] J. Siegrist,[21] G. Signorelli,[31] A. Sill,[38] P. Sinervo,[23] P. Singh,[16] A. J. Slaughter,[44] K. Sliwa,[41] C. Smith,[17] F. D. Snider,[10] A. Solodsky,[35] J. Spalding,[10] T. Speer,[13] P. Sphicas,[22] F. Spinella,[31] M. Spiropulu,[14] L. Spiegel,[10] L. Stanco,[29] J. Steele,[43] A. Stefanini,[31] J. Strologas,[16] F. Strumia, [13] D. Stuart,[10] K. Sumorok,[22] T. Suzuki,[40] R. Takashima,[15] K. Takikawa,[40] M. Tanaka,[40] T. Takano,[28] B. Tannenbaum,[5] W. Taylor,[23] M. Tecchio,[24] P. K. Teng,[1] K. Terashi,[40] S. Tether,[22] D. Theriot,[10] R. Thurman-Keup,[2] P. Tipton,[34] S. Tkaczyk,[10] K. Tollefson,[34] A. Tollestrup,[10] H. Toyoda,[28] W. Trischuk,[23] J. F. de Troconiz,[14] S. Truitt,[24] J. Tseng,[22] N. Turini,[31] F. Ukegawa,[40] J. Valls,[36] S. Vejcik III,[10] G. Velev,[31] R. Vidal,[10] R. Vilar,[6] I. Vologouev,[21] D. Vucinic,[22] R. G. Wagner,[2] R. L. Wagner,[10] J. Wahl,[7] N. B. Wallace,[36] A. M. Walsh,[36] C. Wang,[9] C. H. Wang,[1] M. J. Wang,[1] T. Watanabe,[40] T. Watts,[36] R. Webb,[37] H. Wenzel,[18] W. C. Wester III,[10] A. B. Wicklund,[2] E. Wicklund,[10] H. H. Williams,[30] P. Wilson,[10] B. L. Winer,[27] D. Winn,[24] S. Wolbers,[10] D. Wolinski,[24] J. Wolinski,[25] S. Worm,[26] X. Wu,[13] A. Yagil,[10] W. Yao,[21] G. P. Yeh,[10] P. Yeh,[1] J. Yoh,[10] C. Yosef,[25] T. Yoshida,[28] I. Yu,[19] S. Yu,[30] F. Zetti,[21] and S. Zucchelli[3]

(CDF Collaboration)

[1] *Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China*

[2] *Argonne National Laboratory, Argonne, Illinois 60439*

[3] *Istituto Nazionale di Fisica Nucleare, University of Bologna, I-40127 Bologna, Italy*

[4] *Brandeis University, Waltham, Massachusetts 02254*

[5] *University of California at Los Angeles, Los Angeles, California 90024*

[6] *Instituto de Fisica de Cantabria, University of Cantabria, 39005 Santander, Spain*

[7] *Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637*

[8] *Joint Institute for Nuclear Research, RU-141980 Dubna, Russia*

[9] *Duke University, Durham, North Carolina 27708*

[10] *Fermi National Accelerator Laboratory, Batavia, Illinois 60510*

[11] *University of Florida, Gainesville, Florida 32611*

[12] *Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*

[13] *University of Geneva, CH-1211 Geneva 4, Switzerland*

[14] *Harvard University, Cambridge, Massachusetts 02138*

[15] *Hiroshima University, Higashi-Hiroshima 724, Japan*

[16] *University of Illinois, Urbana, Illinois 61801*

[17] *The Johns Hopkins University, Baltimore, Maryland 21218*

[18] *Institut für Experimentelle Kernphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany*

[19] *Korean Hadron Collider Laboratory: Kyungpook National University, Taegu 702-701; Seoul National University, Seoul 151-742; and SungKyunKwan University, Suwon 440-746; Korea*

[20] *High Energy Accelerator Research Organization (KEK), Tsukuba, Ibaraki 305, Japan*

[21] *Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720*

[22] *Massachusetts Institute of Technology, Cambridge, Massachusetts 02139*

[23] *Institute of Particle Physics: McGill University, Montreal H3A 2T8; and University of Toronto, Toronto M5S 1A7; Canada*

[24] *University of Michigan, Ann Arbor, Michigan 48109*

[25] *Michigan State University, East Lansing, Michigan 48824*

[26] *University of New Mexico, Albuquerque, New Mexico 87131*

[27] *The Ohio State University, Columbus, Ohio 43210*

[28] *Osaka City University, Osaka 588, Japan*

[29] *Universita di Padova, Istituto Nazionale di Fisica Nucleare, Sezione di Padova, I-35131 Padova, Italy*

[30] *University of Pennsylvania, Philadelphia, Pennsylvania 19104*

[31] *Istituto Nazionale di Fisica Nucleare, University and Scuola Normale Superiore of Pisa, I-56100 Pisa, Italy*

[32] *University of Pittsburgh, Pittsburgh, Pennsylvania 15260*

[33] *Purdue University, West Lafayette, Indiana 47907*

[34] *University of Rochester, Rochester, New York 14627*

[35] *Rockefeller University, New York, New York 10021*

[36] *Rutgers University, Piscataway, New Jersey 08855*

[37] *Texas A&M University, College Station, Texas 77843*

[38] *Texas Tech University, Lubbock, Texas 79409*

[39] *Istituto Nazionale di Fisica Nucleare, University of Trieste/ Udine, Italy*

[40] *University of Tsukuba, Tsukuba, Ibaraki 305, Japan*

[41] *Tufts University, Medford, Massachusetts 02155*

[42] *Waseda University, Tokyo 169, Japan*

[43] *University of Wisconsin, Madison, Wisconsin 53706*

[44] *Yale University, New Haven, Connecticut 06520*

Abstract

Top quark production offers the unique opportunity to search for a charged Higgs boson ($H^{\pm}$), as the contribution from $t \to H^{+}b \to \tau^{+}\nu b$ can be large in extensions of the Standard Model. We use results from a search for top quark pair production by the Collider Detector at Fermilab (CDF) in the eτ+$\not\!\!E_T$+jets and $\mu\tau$ + $\not\!\!E_T$+jets signatures to set an upper limit on the branching ratio of $\mathcal{B}(t \to H^{+}b)$. The upper limit is in the range 0.5 to 0.6 at 95% C.L. for H^{+} masses in the range 60 to 160 GeV, assuming the branching ratio for $H^{+} \to \tau\nu$

is 100% . The τ lepton is detected through its 1-prong and 3-prong hadronic decays.

Typeset using REVTEX

Many extensions of the Standard Model (SM) include a Higgs sector with two Higgs doublets, resulting in the existence of charged ($H^{\pm}$) as well as neutral (h, H^0, A) Higgs bosons. The simplest extensions are the Two-Higgs Doublet Models (2HDMs) [1], in which the extension consists only of the extra doublet. In Type I 2HDM models only one of the Higgs doublets couples to fermions, while in Type II models one Higgs doublet couples to the the "up" fermions (e.g., u,c,t), while the other to the "down" fermions (e.g., d,s,b). The Minimal Supersymmetric Model (MSSM) [2] is a further extension of the SM, and has a Higgs sector like that of a Type II 2HDM.

If the charged Higgs boson is lighter than the top quark [3–5], i.e. $m_{H^{\pm}} < (m_{top} - m_b)$, the decay mode $t \to H^+ b$ will compete with the SM decay $t \to W^+ b$. The consequence is that $t\bar{t}$ production and decay will provide a source of Higgs bosons in the channels $W^{\pm}H^{\mp}b\bar{b}$ and $H^+H^-b\bar{b}$ produced with a strong–interaction rather than the weak–interaction cross–section of direct H^+H^- pair production. In addition, the signature from top pair production and decay is much cleaner than that of the direct production with respect to QCD background.

In 2HDM and MSSM models the branching ratio for $t \to H^+ b$, $\mathcal{B}^{\mathrm{t}}_{\mathrm{Hb}}$, depends on the charged Higgs mass and the ratio $\tan\beta$ of the vacuum expectation values for the two Higgs doublets. Figure 1 shows the expected branching ratio from a leading–log QCD calculation [6] in the MSSM for three different charged Higgs masses $m_{H^{\pm}} = 60, 100, 140$ GeV/c^2 as a function of $\tan\beta$. For $\tan\beta \lesssim 1$ and $\tan\beta \gtrsim 70$ the MSSM predicts that the decay mode $t \to H^+ b$ dominates.

Also shown in Figure 1 is the predicted branching ratio in the MSSM at lowest order for the decay of the charged Higgs boson into a charged τ–lepton and a τ–neutrino ($\mathcal{B}^{\mathrm{H}}_{\tau\nu}$), which has little dependence on the charged Higgs mass. For $\tan\beta > 1$ the decay $H^+ \to \tau^+\nu_\tau$ is predicted to dominate over the other main decay mode, $H^+ \to c\bar{s}$, and for $\tan\beta > 5$ the branching ratio $\mathcal{B}^{\mathrm{H}}_{\tau\nu}$ is expected to be nearly 100%. Thus there would be an excess of top events with tau leptons over the number expected from SM events in which $t\bar{t} \to W^+W^-b\bar{b}$, followed by $W \to \tau\nu$.

Recent calculations, however, have shown that at large values of $\tan\beta$ the predicted branching ratio for $t \to H^+ b$ is highly sen-

sitive to higher–order radiative corrections, which are model–dependent [7]. Limits in the $\tan\beta - m_{H^\pm}$ parameter plane consequently depend critically on the parameters of the model. However the direct search for the signature of a τ lepton in top decays allows us to set an upper limit on the branching ratio of $t \to H^+ b$, assuming the branching ratio for $H^+ \to \tau\nu$ is 100%, for example.

Previous searches for the charged Higgs boson in top decay have been in the $\tau + \not{E}_t$ channel [8] [9], $\ell\ell + \not{E}_T$+X (ℓ = e or μ, X=anything) channel [10], the $\not{E}_T + \tau$+jets channel [11,12], the ℓ +jets channel [13,14], and the $\not{E}_T + \tau b + O$+ jet ($O = e$, μ, τ or jet) channel [15]. Both Ref. [11] and Ref. [15] select the events with a $\not{E}_T$ trigger, while Ref. [13,14] is an indirect search using a disappearance method. Searches for direct production at LEP set a lower limit on the mass of 59 GeV/c^2 [16]. Indirect limits have also been set from measurements of the rate for the decay $b \to s\gamma$ [17]. However higher–order calculations have shown that in both 2HDM models [18] and the MSSM [19] these limits are also highly model–dependent.

CDF has published a search for τ leptons from decays of top quark pairs in the $\ell\tau + \not{E}_T$+2 jets+X ($\ell = e, \mu$) channel [20], where the events were selected on the presence of a high–p$_T$ e or μ. We present here the constraints that this analysis (the "$\ell\tau$" analysis) imposes on the branching ratio of the top quark into a charged Higgs boson. This was suggested in Ref. [21], where the authors compare the CDF data with a generator–level Monte Carlo calculation for the number of expected events from charged Higgs decay.

In this paper we start with the number of top candidate events found in the $\ell\tau + \not{E}_T + 2$ jets+X data in the analysis of Ref. [20]. We then apply the same selection criteria to Monte Carlo events that contain top quark pairs in which one or both top quarks decay to the charged Higgs (i.e. $t\bar{t} \to W^\pm H^\mp b\bar{b}$ and $t\bar{t} \to H^+ H^- b\bar{b}$), for different Higgs masses. We assume there are no other top quark decays other than $t \to W^+ b$ and $t \to H^+ b$. We perform a full calculation of the acceptances including detector effects, and determine the expected number of excess events due to Higgs production and subsequent decay. From this we can set a limit on the branching ratio $t \to H^+ b$.

The selection used in this analysis requires high–p_T inclusive lepton events that

contain an electron with $E_T > 20$ GeV or a muon with $p_T > 20$ GeV/c in the central region ($|\eta| < 1.0$). The other lepton must be a tau lepton, also in the central region, with momentum $p_T > 15$ GeV/c [25]. Dilepton events from $t\bar{t}$ decays are expected to contain two jets from b–decays and large missing transverse energy from the neutrinos. Therefore, we select events with ≥ 2 jets (with $E_T > 10$ GeV and $|\eta| < 2.0$), and with large $\not\!\!E_T$ significance ($S_{\not\!\!E_T} > 3$), as described in detail in Ref. [20].

Two complementary techniques, one which identified the τ lepton starting with clusters in the calorimeter, and another which started with a high-P_T single track, were used for identifying hadronically decaying τ's [20]. Here, we combine the two tau selections by accepting events which pass either criteria. Both techniques find the same four top dilepton candidates in 106 pb^{-1} of data. We expect 3.1±0.5 events from background sources, and 0.9 ± 0.1 events from SM top decay. Although the identification of b-quarks was not part of the search criteria, three of the four candidate events contain at least one b–tagged jet [23], which supports the $t\bar{t}$ hypothesis as we expect 0.2 tagged events from background [20]. The total acceptance of the combined selection for SM top quark pairs decays, i.e. the events that pass the final cuts over the number of generated $t\bar{t}$ events, is (0.172±0.014)%. In the following we will use the combined tau selection for our results.

If a charged Higgs boson is present all three of the final states $W^+W^-b\bar{b}$, $W^\pm H^\mp b\bar{b}$, and $H^+H^-b\bar{b}$ can contribute to the the $\ell\tau$ channel. The total acceptance for top decay in the $\ell\tau$ channel is given by:

$$A_{tot}^{\ell\tau} = (1-\mathcal{B}_{\rm Hb}^{\rm t})^2 A_{WW}^{\ell\tau} + 2(1-\mathcal{B}_{\rm Hb}^{\rm t})\mathcal{B}_{\rm Hb}^{\rm t}\mathcal{B}_{\tau\nu}^{\rm H} A_{WH}^{\ell\tau} + (\mathcal{B}_{\rm Hb}^{\rm t})^2(\mathcal{B}_{\tau\nu}^{\rm H})^2 A_{HH}^{\ell\tau} \quad (1)$$

Here $A_{WW}^{\ell\tau}$, is the total acceptance of the event selection criteria for the case where the $t\bar{t}$ pair decays into into $W^+W^-b\bar{b}$. It includes the geometric and kinematic acceptances, the efficiencies for the trigger, lepton identification, and cuts on the event topology, and all branching ratios of both the τ and the W boson [24]. Similarly, $A_{WH}^{\ell\tau}$, and $A_{HH}^{\ell\tau}$ are the respective total acceptances for the $t\bar{t}$ pair decays into $W^\pm H^\mp b\bar{b}$, and $H^+H^-b\bar{b}$, but where the branching ratio of the top to Higgs ($\mathcal{B}_{\rm Hb}^{\rm t}$) and of the Higgs to tau ($\mathcal{B}_{\tau\nu}^{\rm H}$) have been fac-

tored out explicitly. We assume that $\mathcal{B}^{\rm H}_{\tau\nu}$ is 100%, as it would be at large $\tan\beta$ in the MSSM, and set a limit on $\mathcal{B}^{\rm t}_{\rm Hb}$.

We use a top quark mass of 175 GeV/c^2. Monte Carlo simulations of $t\bar{t}$ production and decay in the three modes $W^+W^-b\bar{b}$, $W^\pm H^\mp b\bar{b}$, and $H^+H^-b\bar{b}$ provide estimates of the geometric and kinematic acceptance, $A_{geom\cdot P_T}$, and of the efficiency of the cuts on the event topology for different Higgs masses ($m_{H^\pm} = 60, 80, 100, 120, 140, 160$ GeV/c^2). We use the PYTHIA [22] Monte Carlo to generate $t\bar{t}$ events, the TAUOLA package [26], which correctly treats the τ polarization, to decay the tau lepton, and a detector simulation. The selection of events is identical to that described in detail in Ref. [20]. The efficiency for electron and muon identification is measured from $Z^\circ \to e^+e^-$ and $Z^\circ \to \mu^+\mu^-$ data.

Figure 2 shows $A_{geom\cdot P_T}$, the efficiency ϵ_{jet} of the 2–jet cut, the efficiency ϵ_{H_T} of the cut on the total transverse energy H_T, and the efficiency of the cut on the $\not\!\!E_T$ significance, as a function of Higgs mass. As $m_{H^\pm}$ increases the tau leptons become more energetic and $A_{geom\cdot P_T}$ increases. When $m_{H^\pm}$ approaches m_{top} the b–jets instead become less energetic and ϵ_{jet} drops rapidly. Figure 3 shows the resulting values for $A^{\ell\tau}_{WH}$ and $A^{\ell\tau}_{HH}$ versus $m_{H^\pm}$; the numerical values are listed in Table I. Note that relative to A_{HH}, A_{WH} has a factor of 2/9 in it due to the branching ratio for $W \to \ell\nu$, and a factor of 2 due to the two possible charge combinations W^+H^- and W^-H^+. Overall, the total acceptance ($A^{\ell\tau}_{tot}$) is rather insensitive to the value of the Higgs mass, being about 0.7–1.3 % until $m_{H^\pm}$ approaches m_{top}. This is to be compared to the acceptance $A^{l\tau}_{WW}$ in the W^+W^- final state [20] of 0.17%.

The expected number of events in the $\ell\tau$ channel is given by:

$$N^{\ell\tau}_{exp} = \sigma_{t\bar{t}} \cdot \mathcal{L} \cdot A^{\ell\tau}_{tot}(\mathcal{B}^{\rm t}_{\rm Hb}, m_{H^\pm}) \qquad (2)$$

and depends on $\mathcal{B}^{\rm t}_{\rm Hb}$, the Higgs boson mass, and $\sigma_{t\bar{t}}$, the total top pair production cross section. Rather than use the theoretical prediction for $\sigma_{t\bar{t}}$, we normalize to the observed number of events in the 'lepton + jets' channel with a secondary vertex tag, taking into account the contributions from the three separate decay final states of $W^+W^-b\bar{b}$, $W^\pm H^\mp b\bar{b}$, and $H^+H^-b\bar{b}$. We use the full Monte Carlo simulation and the updated

tagging efficiency [27], which gives a central value of $\sigma_{t\bar{t}} = 5.1$ pb in the SM case of $\mathcal{B}^{\rm t}_{\rm Hb} = 0$. We thus calculate $\sigma_{t\bar{t}}$ from the number of observed events in the lepton plus jets channel with a secondary vertex tag, $N^{\ell+jets}$, the expected number of SM background events $B^{\ell+jets}$, and a total acceptance $A^{\ell+jets}_{tot}(\tan\beta, m_{H^\pm})$ that takes the $W^\pm H^\mp b\bar{b}$ and $H^+H^-b\bar{b}$ decay modes into account:

$$\sigma_{t\bar{t}} = \frac{N^{\ell+jets} - B^{\ell+jets}}{\mathcal{L} \cdot A^{\ell+jets}_{tot}(\mathcal{B}^{\rm t}_{\rm Hb}, m_{H^\pm})} \qquad (3)$$

where $A^{\ell+jets}_{tot}$ is given analogously to $A^{\ell\tau}_{tot}$ by:

$$\begin{aligned} A^{\ell+jets}_{tot} = {} & (1-\mathcal{B}^{\rm t}_{\rm Hb}))^2 A^{\ell+jets}_{WW} + \\ & 2(1-\mathcal{B}^{\rm t}_{\rm Hb})\mathcal{B}^{\rm t}_{\rm Hb} A^{\ell+jets}_{WH} + \\ & (\mathcal{B}^{\rm t}_{\rm Hb})^2 A^{\ell+jets}_{HH} \end{aligned} \qquad (4)$$

The contribution from $H^+ \to c\bar{s}$ decays is neglected, as we have assumed $\mathcal{B}^{\rm H}_{\tau\nu} = 1$. For a large branching ratio into H^+b, the $H^+H^-b\bar{b}$ mode becomes dominant and the leptons (e or μ), which in this case originate from tau decays, have a softer p_T spectrum than leptons produced in W decays, and $A^{\ell+jets}_{tot}$ decreases. Figure 4 shows the expected number of events versus $\mathcal{B}^{\rm t}_{\rm Hb}$ from each of the $W^+W^-b\bar{b}$, $W^\pm H^\mp b\bar{b}$, and $H^+H^-b\bar{b}$ decay modes for $m_{H^\pm} = 100$ GeV/c^2.

Based on the observation of 4 events and the predicted background of 3.1 ± 0.5 events, we calculate a 95% C.L. upper limit on Higgs production of 7.4 events. When calculating the limit, we include the systematic uncertainties, which are dominated by uncertainties on $\sigma_{t\bar{t}}$ (19%), tau identification (11%), b–tagging efficiency (10%) and Monte Carlo statistics (8%). Then, to determine a limit on the branching ratio $\mathcal{B}^{\rm t}_{\rm Hb}$, we calculate the number of events expected versus $\mathcal{B}^{\rm t}_{\rm Hb}$ for different Higgs masses in steps of 20 GeV/c^2, assuming the branching ratio for $H^+ \to \tau\nu$ is 100%. Figure 5 shows the region excluded at 95% C.L. as a function of the branching ratio of $t \to H^+b$. The upper limit is in the range 0.5 to 0.6 at 95% C.L. for H^+ masses in the range 60 to 160 GeV.

For the special case of the MSSM, although the branching ratios have been shown to be strongly model–dependent, for $\mu < 0$ the SUSY QCD and QCD corrections come close to cancelling, and the prediction is almost unchanged from the tree–level prediction [7]. Figure 6 shows the expected number of $\ell\tau$ events versus $\tan\beta$ from each of the $W^+W^-b\bar{b}$, $W^\pm H^\mp b\bar{b}$, and $H^+H^-b\bar{b}$ decay modes for $m_{H^\pm} = 100$ GeV/c^2, at lowest order in the MSSM. The shapes of the

curves are mainly due to the variation of the BR($t \to H^+ b$) as a function of $\tan\beta$. Figure 7 shows the excluded region in the plane of $m_{H^\pm}$ and $\tan\beta$, again at lowest order in the MSSM. In the region at large values of $\tan\beta$ the tbH^+ Yukawa coupling may become non–perturbative (see Ref. [7]). In this case the limit is not valid.

We compare our results to those of Ref. [21]. We find that the acceptance is smaller by about a factor of two. The limits presented in this letter use the correct $W^+W^-b\bar{b}$, $W^\pm H^\mp b\bar{b}$, and $H^+H^-b\bar{b}$ acceptances, including the correlations among the different objects (e, μ, τ, b-quark) in the events. The insight of Ref [21] that this will be a channel of much interest in Fermilab Run II remains intact, however.

In conclusion, we have used the data from the CDF search [20] for top quark decays into final states containing a light lepton (e or μ) and a τ lepton, detected through its 1-prong and 3-prong hadronic decays, to set a limit on the branching ratio of the top quark into the charged Higgs plus a b quark, $\mathcal{B}^{\rm t}_{\rm Hb}$. The limit ranges from 0.5 to 0.6 at 95% C.L. for H^+ masses in the range 60 to 160 GeV, assuming the branching ratio for $H^+ \to \tau\nu$ is 100%.

We thank D.P. Roy for stimulating our interest in this analysis and for discussions, and G. Farrar for (long–ago) suggesting that we use the $\mathcal{B}^{\rm t}_{\rm Hb} - m_{H^\pm}$ plane rather than the $\tan\beta - m_{H^\pm}$ plane for presenting limits. We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Science and Culture of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China, and the A.P. Sloan Foundation.

REFERENCES

[1] For an introduction, see J. Gunion, H. Haber, G. Kane, and S. Dawson, *The Higgs Hunter's Guide*, Frontiers in Physics, Addison Wesley, 1990. See also Ref. [7]

[2] For an introduction, see Martin, S.P., *A Supersymmetry Primer*, in *Perspectives on Supersymmetry*, G. Kane ed., World Scientific, 1998

[3] F. Abe *et al.* (The CDF Collaboration), *Phys. Rev.* D **50**, 2966 (1994)

[4] F. Abe *et al.* (The CDF Collaboration), Phys. Rev. Lett. **74**, 2626 (1995)

[5] S. Abachi *et al.* (The DØ Collaboration), Phys. Rev. Lett. **74**, 2632 (1995)

[6] M. Drees and D.P. Roy, Phys. Lett. B **269**, 155 (1991); D.P. Roy, *ibid.* **283**, 403 (1992)

[7] J.A. Coarasa, J. Guasch, J. Sola, Preprint UAB–FT–451, Feb. 1999, hep–ph/9903212

[8] In the CDF coordinate system, θ and ϕ are the polar and azimuthal angles, respectively, with respect to the proton beam direction. The pseudorapidity η is defined as $-\ln\tan(\theta/2)$. The transverse momentum of a particle is $P_T = P\sin\theta$. The analogous quantity using energies, defined as $E_T = E\sin\theta$, is called transverse energy. The missing transverse energy $\not\!\!E_T$ is defined as $-\sum E_T^i\ \hat{r}_i$, where $\hat{r}_i$ are the unit vectors in the transverse plane pointing to the energy depositions in the calorimeter.

[9] C. Albajar *et al.* (The UA1 Collaboration), Phys. Lett. **B257**, 459 (1991); J. Alitti *et al.* (The UA2 Collaboration), Phys. Lett. B **280**, 137 (1992)

[10] F. Abe *et al.* (The CDF Collaboration), Phys. Rev. Lett. **73**, 2667 (1994);

[11] F. Abe *et al.* (The CDF Collaboration), Phys. Rev. Lett. **72**, 1977 (1994)

[12] F. Abe *et al.* (The CDF Collaboration), *Phys. Rev.* D **54**, 735 (1996)

[13] B. Abbott *et al.* (The D0 Collaboration), Phys. Rev. Lett. **82**, 4975 (1999)

[14] B. Bevensee (The CDF Collaboration), Proc. of the 33rd Rencontres de Moriond, QCD and High Energy Hadronic Inter-

actions, 21–28 March 1998, Les Arcs, France; FERMILAB–CONF–98/155–E

[15] F. Abe *et al.* (The CDF Collaboration), Phys. Rev. Lett. **79**, 357 (1997)

[16] R. Barate *et al.* (The Aleph Collaboration) submitted to Phys.Lett.B, hep–ex/9902031; P. Abreu et al. (The DELPHI Collaboration), Phys. Lett. B460, 484 (1999); M. Acciarri et al. (The L3 Collaboration), Phys. Lett. B446, 368 (1999); K. Ackerstaff et al. (The OPAL Collaboration) Phys. Lett. B426, 180 (1998)

[17] M. Alam *et al.* (The CLEO Collaboration), Phys. Rev. Lett 74, 2885 (1995)

[18] F. Borzumati, and C. Greub, Phys. Rev. D58:074004, 1998; Phys. Rev. D59:057501 (1999).

[19] J. Coarasa, J. Guasch, W. Hollik, and J. Sola, Phys. Lett. **B442**, 326 (1998)

[20] F. Abe *et al.* (The CDF Collaboration), Phys. Rev. Lett. **79**, 3585 (1997); M. Gallinaro, Ph.D thesis University of Rome, 1996; M. Hohlmann, Ph.D thesis, University of Chicago, 1997.

[21] M. Guchait and D.P. Roy, Phys. Rev. D **55**, 7263 (1997)

[22] T. Sjöstrand, Comput. Phys. Commun. **82**, 74 (1994)

[23] This is using the same SVX + SLT b-tagging algorithms used in Ref [3]. We emphasize again that in Ref [20] (and consequently in this paper) we did not use b-tagging in the event selection, because of the small number of events expected from top decays.

[24] The reason the acceptance is quoted as containing the branching ratios is that $t\bar{t}$ events are complex, and can satisfy the event selection criteria via a number of different paths, such as the e or μ coming from the W and the τ from the H in the $WHb\bar{b}$ intermediate state, or, the τ coming from the W decay and the e or μ coming from the decay of a τ from the H, for example. We consequently run the Monte Carlo with all decay modes turned on except those involving the Higgs, for which the branching ratios are varied explicitly to find the acceptance for each value. The acceptance is then defined as

the number of accepted events over the number of $t\bar{t}$ events generated.

[25] The selection criteria are that either a track exist with $P_T > 15$ GeV/c or the sum of the transverse momenta of the tracks and the transverse energies of π^0's be greater than 15 GeV/c. See Ref [20] for details.

[26] S. Jadach *et al.*, TAUOLA 2.5, CERN preprint CERN–TH–6793 (1992)

[27] F. Ptohos, Proceedings of the European Physical Society, Finland 1999

TABLES

M_{Higgs}	$A^{l\tau}_{WH}$ (%)	$A^{l\tau}_{HH}$ (%)
60	0.91±0.06	1.00±0.06
80	0.98±0.06	1.17±0.06
100	1.11±0.06	1.32±0.07
120	1.08±0.06	1.32±0.07
140	0.67±0.05	0.98±0.06
160	0.72±0.05	0.32±0.03

TABLE I. The total acceptance versus the mass of the charged Higgs boson for the $\ell\tau + \not{E}_T + 2$ jets $+X$ analysis. The uncertainties are statistical only. These numbers are to be compared to the acceptance for SM top quark pair decays of $A^{l\tau}_{WW} = (0.172 \pm 0.014)\%$. The larger acceptance with the charged Higgs is primarily due to the larger branching fractions into τ leptons.

FIGURES



FIG. 1. Branching fraction of $H \to \tau\nu$ and $t \to Hb$ as a function of $\tan\beta$ at lowest order in the MSSM. The top quark mass is taken as 175 GeV/c^2.



FIG. 2. Contributions to the total acceptance in the $\ell\tau + \not{E} + 2$ jets$+X$ channel versus the mass of the charged Higgs boson. The circles are for the $W^{\pm}H^{\mp}b\bar{b}$ decay of the $t\bar{t}$ pair; the squares for for the $H^+H^-b\bar{b}$ decay. The branching ratio for $H \to \tau\nu$ is assumed to be 100%.



FIG. 3. Total acceptance in the "tau dilepton" channel, versus the mass of the Higgs boson. The circles are for the $W^{\pm}H^{\mp}b\bar{b}$ decay of the $t\bar{t}$ pair; the squares for for the $H^+H^-b\bar{b}$ decay.



FIG. 4. The predicted number of events for 106 pb^{-1} of data versus the branching ratio for top decay into H^+b for M_{Higgs}=100 GeV/c^2. The graph shows the contributions from the $W^+W^-b\bar{b}$, $W^{\pm}H^{\mp}b\bar{b}$, and $H^+H^-b\bar{b}$ channels separately.



FIG. 5. The region excluded at 95% C.L. for charged Higgs production versus the branching ratio for top decay into H^+b.



FIG. 6. The predicted number of events at lowest order in the MSSM for 106 pb^{-1} of data versus $\tan\beta$, for M_{Higgs}=100 GeV/c^2. The graph shows the different contributions from the $H^+H^-b\bar{b}$ and $W^{\pm}H^{\mp}b\bar{b}$ channels separately.



FIG. 7. Excluded regions (95% C.L.) at different values of $\tan\beta$ for charged Higgs production, at lowest order in the MSSM. The coupling tbH^+ may become non–perturbative in the region at large values of $\tan\beta$, and the limit does not apply.